For Research Use Only

SAMSN1 Polyclonal antibody

Catalog Number: 13063-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number: GenBank Accession Number: 13063-1-AP BC029112
Size: GeneID (NCBI): 64092
Source: UNIPROT ID: Rabbit Q9NSI8

Full Name:

SAM domain, SH3 domain and nuclear

Immunogen Catalog Number: localization signals 1

AG3755 Calculated MW:

42 kDa
Observed MW:
42-50 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC

Isotype:

Species Specificity: human, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:20-1:200

Positive Controls:

WB: K-562 cells, IP: K-562 cells,

IHC: human gliomas tissue,
IF/ICC: HepG2 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Satoshi Sueoka	25805236	Ann Surg Oncol	IHC
Mitsuro Kanda	28105178	Oncol Lett	IHC
Zoe Schaefer	39366656	Mol Cell Proteomics	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13063-1-AP (SAMSN1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13063-1-AP (SAMSN1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-SAMSN1 (IP:13063-1-AP, 4ug: Detection:13063-1-AP 1:1000) with K-562 cells lysate 4000 ug.



K-562 cells were subjected to SDS PAGE followed by western blot with 13063-1-AP (SAMSN1 Antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells using 13063-1-AP (SAMSN1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit lgG(H+L).