## For Research Use Only

## SMARCA5 Polyclonal antibody

Catalog Number:13066-1-AP

Featured Product 5 Publications

Antibodies | ELISA kits | Proteins www.ptglab.com


Basic Information	Catalog Number: 13066-1-AP	GenBank Accession Number: BC023144	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	200 ug/ml	8467	WB 1:500-1:2000	
	Source: Rabbit Isotype:	UNIPROT ID: 060264	IHC 1:500-1:2000 IF/ICC 1:500-1:2000	
		Full Name:	0204	
	IgG SWI/SNF related, matrix associated,		ted,	
	Immunogen Catalog Number:	actin dependent regulator of chromatin, subfamily a, member 5 Calculated MW: 1052 aa, 122 kDa		
	AG3759			
				Observed MW:
			122 kDa	
	Applications	Tested Applications:	Positive Controls:	
WB, IHC, IF/ICC, ELISA		WB : HeLa	WB : HeLa cells,	
Cited Applications: WB, IF, IP		IHC : human stomach tissue, IF/ICC : HeLa cells,		
Species Specificity: human				
Cited Species: human, mouse				
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
	retrieval may be performed			
Background Information	retrieval may be performed buffer pH 6.0 SMARCA5, also named as SNF2H overexpressed in CD34-positive e most conserved chromatin remod respect to the histone octamer to	<b>d with citrate</b> or WCRF135, is a 1052 amino acid pro crythrocyte progenitor cells in acute m eling factors. It exists in a variety of o	ligosubunit complexes that move DNA w al arrays. Smarca5 interacts with differen	
	retrieval may be performed buffer pH 6.0 SMARCA5, also named as SNF2H overexpressed in CD34-positive e most conserved chromatin remod respect to the histone octamer to accessory proteins and represents	d with citrate or WCRF 135, is a 1052 amino acid pro erythrocyte progenitor cells in acute m eling factors. It exists in a variety of o generate regularly spaced nucleosom	nyeloid leukemia. SMARCA5 is one of the Iligosubunit complexes that move DNA w al arrays. Smarca5 interacts with differen	
	retrieval may be performed buffer pH 6.0 SMARCA5, also named as SNF2H overexpressed in CD34-positive e most conserved chromatin remod respect to the histone octamer to accessory proteins and represents Author	d with citrate or WCRF135, is a 1052 amino acid pro crythrocyte progenitor cells in acute m eling factors. It exists in a variety of o generate regularly spaced nucleosom a molecular motor for DNA replicatio	nyeloid leukemia. SMARCA5 is one of the ligosubunit complexes that move DNA w al arrays. Smarca5 interacts with differen n, repair, and transcription.	
Background Information	retrieval may be performed buffer pH 6.0 SMARCA5, also named as SNF2H overexpressed in CD34-positive e most conserved chromatin remod respect to the histone octamer to accessory proteins and represents Author Haiyan An	or WCRF 135, is a 1052 amino acid pro erythrocyte progenitor cells in acute m eling factors. It exists in a variety of o generate regularly spaced nucleosom a molecular motor for DNA replicatio Pubmed ID Journal	nyeloid leukemia. SMARCA5 is one of the ligosubunit complexes that move DNA w al arrays. Smarca5 interacts with differen n, repair, and transcription. Application	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





HeLa cells were subjected to SDS PAGE followed by western blot with 13066-1-AP (SMARCAS Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunofluorescent analysis of (4% PFA) fixed HeLa cells using SMARCA5 antibody (13066-1-AP) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 13066-1-AP (SMARCA5 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).