For Research Use Only

NKX3-1 Polyclonal antibody

Catalog Number:13069-1-AP 2 Publications



Basic Information

Catalog Number: 13069-1-AP

Size:
350 µg/ml
Source:
Rabbit
Isotype:

Immunogen Catalog Number:

AG4070

Observed MW: 34-38 kDa

BC074864

4824

Q99801 Full Name:

GeneID (NCBI):

UNIPROT ID:

NK3 homeobox 1
Calculated MW:

234 aa, 26 kDa

GenBank Accession Number:

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000 IHC 1:400-1:1600

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

IHC

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, human testis tissue IHC: human prostate cancer tissue,

Background Information

NKX3-1, also named as NKX3.1 and NKX3A, belongs to the NK-3 homeobox family. It is a transcription factor, which binds preferentially the consensus sequence 5'-TAAGT[AG]-3' and can behave as a transcriptional repressor. NKX3-1 plays an important role in normal prostate development, regulating proliferation of glandular epithelium and in the formation of ducts in prostate. It act as a tumor suppressor controlling prostate carcinogenesis, as shown by the ability to inhibit proliferation and invasion activities of PC-3 prostate cancer cells(PMID:19462257). The calcualted molecular weight of NKX3-1 is 26 kDa, but modified NKX3-1 is about 32-35 kDa. (PMID: 18597829)

Notable Publications

Author	Pubmed ID	Journal	Application
Han Zhang	39638850	Sci Rep	IHC
Weiran Feng	38585869	bioRxiv	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 13069-1-AP (NKX3-1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



LNCaP cells were subjected to SDS PAGE followed by western blot with 13069-1-AP (NK/3-1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 13069-1-AP (NKX3-1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 13069-1-AP (NKX3-1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).