## For Research Use Only

# **UGDH** Polyclonal antibody

Catalog Number: 13151-1-AP

Featured Product

5 Publications



**Basic Information** 

Catalog Number:

13151-1-AP

Size:

450 µg/ml

Source:

Rabbit

GenBank Accession Number:

BC022781

GeneID (NCBI):

7358

Surce:

UNIPROT ID:

Rabbit

O60701

Isotype:

Full Name:

gG UDP-glucose dehydrogenase

Immunogen Catalog Number: Calculated MW: AG3767 494 aa, 55 kDa

Observed MW: 55-60 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, HepG2 cells, mouse liver tissue, rat

**Purification Method:** 

WB 1:500-1:2000 IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

liver tissue

IHC: human liver cancer tissue,

IF/ICC: A549 cells,

# **Background Information**

UDP-glucose 6-dehydrogenase (UGDH) is a key enzyme in the uronic acid pathway, and converts UDP-glucose (UDP-Glc) to UDP-glucuronic acid (UDP-GlcA). UGDH is critical to the production of extracellular matrix components which are essential to the migration and connectivity of neurons early in human brain development. UDP-GlcA is an essential sugar nucleotide precursor and a key component in the synthesis and stepwise degradation of glycosaminoglycans (GAGs). The protein subunit migrates on SDS-PAGE at ~55 kDa. (PMID: 32175296, PMID: 21576248, PMID: 31243371).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yinxian Wen	25202993	PLoS One	WB, IHC
Qi Ying	25974208	PLoS One	WB
Xiongjun Wang	31243371	Nature	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 13151-1-AP (UGDH antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 13151-1-AP (UGDH antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A549 cells using UGDH antibody (13151-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).