

For Research Use Only

TBX4/5 Polyclonal antibody

Catalog Number: 13178-1-AP **11 Publications**



Basic Information

Catalog Number: 13178-1-AP	GenBank Accession Number: BC027942	Purification Method: Antigen affinity purification
Concentration: 273 µg/ml	GeneID (NCBI): 6910	Recommended Dilutions: WB 1:500-1:1000 IF/ICC 1:10-1:100
Source: Rabbit	UNIPROT ID: Q99593	
Isotype: IgG	Full Name: T-box 5	
Immunogen Catalog Number: AG3845	Calculated MW: 518 aa, 57 kDa	
	Observed MW: 58 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells, MCF-7 cells IF/ICC : HepG2 cells,
Cited Applications: WB, IF	
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Background Information

T-box genes encode a family of transcription factors that have been identified in all metazoans and which play diverse roles during embryonic development. TBX5, a member of T-box family, has an essential role in the initiation of limb outgrowth and heart development. Genetic variation near TBX5 that associates with PR and/or QRS interval variation, which suggests that TBX5 plays a role in CCS function in the general population. Because 77 % amino acid sequence of TBX5 is identical to TBX4, this antibody recognizes both TBX4 and TBX5.

Notable Publications

Author	Pubmed ID	Journal	Application
De-Gang Wang	34490705	Am J Med Genet A	IF
Zhentaο Zhang	31628386	Sci Rep	WB
Zhihua Ren	35609006	Environ Sci Technol	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 13178-1-AP (TBX5 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells using 13178-1-AP (TBX5 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).