For Research Use Only

GBP5 Polyclonal antibody

Catalog Number: 13220-1-AP

Featured Product

44 Publications



Basic Information

Catalog Number:

13220-1-AP

BC031639

Concentration:

650 ug/ml

115362

Source:

UNIPROT ID:

Rabbit

Q96PP8

Isotype:

GenBank Accession Number:

BC031639

GeneID (NCBI):

115362

UNIPROT ID:

Q96PP8

Full Name:

gG guanylate binding protein 5

Immunogen Catalog Number:Calculated MW:AG4103586 aa, 67 kDa

Observed MW: 65 kDa, 55 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IHC 1:250-1:1000

WB 1:2000-1:16000

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC

Species Specificity: human, rat Cited Species: human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: U-937 cells, Jurkat cells

IP: U-937 cells,
IHC: rat spleen tissue,

Background Information

Guanylate binding protein (GBP) 5 belongs to the GBP family, which is involved in important cellular processes, including signal transduction, translation, vesicle trafficking, and exocytosis. GBP5 was reported to be a critical cellular factor in inflammasome assembly. Furthermore, GBP5 has been reported to induce inflammatory cytokines, including IL-1 beta and IL-18, through the activation of NLRP3 and AIM2. GBP5 encodes two isoforms: GBP-5a/b (67 kDa) and GBP-5ta (55 kDa)

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|---------------------|-------------|
| Naiquan Liu | 36175170 | Immunol Invest | WB,IHC |
| Ting-Ting Yu | 34562845 | Int Immunopharmacol | WB |
| Hua Zhen Xu | 34488792 | J Nanobiotechnology | WB,IHC |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




U-937 cells were subjected to SDS PAGE followed by western blot with 13220-1-AP (GBP5 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




IP result of anti-GBP5 (IP:13220-1-AP, 4ug; Detection:13220-1-AP 1:1000) with U-937 cells lysate 4000 ug.



Immunohistochemical analysis of paraffinembedded rat spleen tissue slide using 13220-1-AP (GBP5 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat spleen tissue slide using 13220-1-AP (GBP5 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 MCF-7 cells were intracellularly stained with 0.4 ug GBP5 Polyclonal antibody (13220-1-AP) and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).