For Research Use Only

ASPA Polyclonal antibody

Catalog Number: 13244-1-AP

Featured Product

5 Publications

BC029128

GeneID (NCBI):

UNIPROT ID:

P45381 Full Name:

GenBank Accession Number:



Purification Method:

WB 1:1000-1:6000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 13244-1-AP Size:

Isotype:

500 µg/ml Source: Rabbit

IgG aspartoacylase (Canavan disease)

Immunogen Catalog Number:Calculated MW:AG4029313 aa, 36 kDaObserved MW:

36 kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse brain tissue,

IHC: human gliomas tissue,

Background Information

ASPA(Aspartoacylase) also called aminoacylase-2, is an enzyme that hydrolyzes N-acetyl-L-aspartic acid (NAA) to aspartate and acetate. Human ASPA encodes a deduced 313-amino acid protein with a molecular mass of 36 KDa. It shares 92% sequence identity with the bovine protein and contains 1 potential N-glycosylation site and 5 phosphorylation sites. Defects in ASPA are the cause of Canavan disease (CAND). This protein can exsit as a homodimer (18293939).

Notable Publications

Author	Pubmed ID	Journal	Application
Vasiliki-Ilya Gargareta	35543322	Elife	WB
Kaiyun Yao	35281914	Front Pharmacol	WB
Sophie B Siems	39344832	Glia	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data







Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13244-1-AP (ASPA antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 13244-1-AP (ASPA antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.