For Research Use Only

TEAD1 Polyclonal antibody

Catalog Number: 13283-1-AP

Featured Product

12 Publications



Purification Method:

WB 1:500-1:1500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 13283-1-AP Concentration: 800 µg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG4082

GenBank Accession Number:

BC026959 GeneID (NCBI): 7003

UNIPROT ID:

P28347

52 kDa

Full Name: TEA domain family member 1 (SV40 transcriptional enhancer factor)

Calculated MW: 426 aa, 47, 9 kDa Observed MW:

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IF

Species Specificity: mouse, human Cited Species: human, rat Positive Controls:

WB: BGC-823 cells, A375 cells, C2C12 cells

Background Information

TEAD1 belongs to a protein family that share the TEA/ATTS domain. It acts as a transcriptional factor by binding to the GT-IIC and Sph(I 1 II) enhansons in the simian virus 40 (SV40) enhancer. Also it can modulate SV40 late transcription together with large T antigen. The transcriptional activity of TEAD1 required several regions of the proteins, a TBP-associated factors and a limiting transcriptional intermediary factors. TEAD1 also preforms an essential role in the Hippo signaling pathway. TEAD1 regulates gene expression of YAP1 and WWTR1/TAZ, thus mediating cell proliferation, migration and epithelial mesenchymal transition induction. TEAD is preferentially expressed in skeletal muscle. Lower levels in pancreas, placenta, and heart.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------------|-----------|---------------------|-------------|
| Chunli Lin | 28892790 | Biomed Pharmacother | WB |
| Jian-Jia Cao | 25175178 | Oncol Rep | WB |
| Rina Fujiwara-Tani | 32481659 | Int J Mol Sci | WB |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



BGC-823 cells were subjected to SDS PAGE followed by western blot with 13283-1-AP (TEAD1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.