For Research Use Only

BUB1 Polyclonal antibody

Catalog Number: 13330-1-AP

Featured Product

11 Publications



Purification Method:

WB 1:500-1:1000 IF/ICC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

Basic Information

Applications

Catalog Number: 13330-1-AP Size: 450 µg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG4138

Tested Applications:

WB, IF/ICC, ELISA Cited Applications: WB, IHC, IF

Species Specificity: human, canine **Cited Species:**

human, carassius auratus

GenBank Accession Number:

BC028201 GeneID (NCBI):

UNIPROT ID: 043683

Full Name:

budding uninhibited by

benzimidazoles 1 homolog (yeast)

Calculated MW: 1085 aa, 122 kDa

Observed MW: 45 kDa, 120 kDa

Positive Controls:

WB: K-562 cells, HeLa cells IF/ICC: IMR-90 cells, MDCK cells

Background Information

Bub1 is an evolutionarily conserved mitotic checkpoint control protein that is present in diverse organisms including yeast and humans. Bub1 is a serine/threonine protein kinase and is required for recruitment of Mad1, Mad2, Bub3, and CENP-E to kinetochores. Full-length Bub1 protein was cleaved to an 45 kDa fragment and an intermediate fragment of 90 kDa, only detected at early stages of apoptosis execution, indicating cleavage of Bub1 in the amino- and carboxy-terminal regions of the protein(PMID:15818396).

Notable Publications

Author	Pubmed ID	Journal	Application
Xie Mengyan	36176446	Front Pharmacol	WB
Junjie Piao	30898709	Gene	IHC
Liang Chen	35659274	J Exp Clin Cancer Res	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



IF results of BUB1 anitbody with IMR-90 cells by M.S. Grasiella Angelina Andriani Cristina Montagna Lab at Albert Einstein College of Medicine-Bronx -NY.



MKN-45 cells were subjected to SDS PAGE followed by western blot with 13330-1-AP (BUB1 antibody) at dilution of 1:300 incubated at room temperature for 15 back.



K-562 cells were subjected to SDS PAGE followed by western blot with 13330-1-AP (BUB1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.