For Research Use Only

RAD17 Polyclonal antibody

Catalog Number: 13358-1-AP

Featured Product

3 Publications



Purification Method:

WB: 1:1000-1:6000 IHC: 1:20-1:200

IF/ICC: 1:50-1:500

100 µl suspension

Antigen affinity purification

FC (Intra): 0.25 ug per 10^6 cells in a

Recommended Dilutions:

Basic Information

Catalog Number:

13358-1-AP

BC032304

Concentration:

600 ug/ml

5884

Source:

UNIPROT ID:

Rabbit

O75943

Isotype:

GenBank Accession Number:

BC032304

GeneID (NCBI):

5884

UNIPROT ID:

75943

Full Name:

gG RAD17 homolog (S. pombe)

Immunogen Catalog Number:Calculated MW:AG4171681 aa, 71 kDaObserved MW:

71 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB

Species Specificity:

human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: K-562 cells, Raji cells

IHC: human breast cancer tissue,

IF/ICC : HeLa cells,
FC (Intra) : A431 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Jill E Hunter	36240066	Biochem J	WB
Jill E Hunter	36240065	Biochem J	WB
Emily M Schleicher	35717336	Oncogenesis	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13358-1-AP (RAD17 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer using 13358-1-AP (RAD17 antibody) at dilution of 1:100 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human breast cancer using 13358-1-AP (RAD17 antibody) at dilution of 1:100 (under 40x lans)



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 13358-1-AP (RAD17 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1x10^6 A431 cells were intracellularly stained with 0.25 ug RAD17 Polyclonal antibody (13358-1-AP) and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).