For Research Use Only

SPA17 Polyclonal antibody

Catalog Number: 13367-1-AP

5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 13367-1-AP BC032457 GeneID (NCBI): Size: 750 µg/ml 53340 **UNIPROT ID:** Source: Rabbit Q15506 Full Name: Isotype:

sperm autoantigenic protein 17

Calculated MW: Immunogen Catalog Number: AG4186 151 aa, 17 kDa Observed MW:

25 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2400

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse testis tissue, IP: mouse testis tissue,

IHC: human testis tissue, human ovary tumor tissue

Background Information

Sperm surface protein Sp17(SPA17)is a 151 amino acids protein (17kDa), which is a sperm surface peripheral membrane protein. It is a highly conserved mammalian protein expressed in developing spermatozoa and mature spermatids, and reported to play a role in the interaction of sperm with the zona pellucida. SP17 exists as a $homodimer and \ localizes \ to \ the \ head \ and \ tail \ of \ spermatozoa. \ SP17 \ has \ been \ identified \ as \ a \ cancer/test \ is \ antigen$ and is expressed in ovarian cancer and multiple myeloma. This suggests that SP17 could be suitable as a target in tumor immunotherapy.

Notable Publications

Author	Pubmed ID	Journal	Application
Takujiro Homma	33087733	Sci Rep	WB
Laurie L Brunette	30309325	BMC Cancer	IHC
Zewei Tu	35592314	Front Immunol	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




mouse testis tissue were subjected to SDS PAGE followed by western blot with 13367-1-AP (SPA17 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 13367-1-AP (SPA17 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 13367-1-AP (SPA17 antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-SPA17 (IP:13367-1-AP, 3ug; Detection:13367-1-AP 1:400) with mouse testis tissue lysate 4800ug.