

For Research Use Only

TDG Polyclonal antibody

Catalog Number: 13370-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number:

13370-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4190

GenBank Accession Number:

BC037557

GeneID (NCBI):

6996

UNIPROT ID:

Q13569

Full Name:

thymine-DNA glycosylase

Calculated MW:

410 aa, 46 kDa

Observed MW:

55-60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, IP, WB, ELISA

Cited Applications:

WB, IP, IHC, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB: Caco-2 cells, mouse colon tissue, HeLa cells, human brain tissue, Raji cells, mouse thymus tissue, HCT-116 cells, Jurkat cells, K-562 cells, MCF-7 cells

IP: U-937 cells,

IF: HeLa cells,

Background Information

TDG belongs to the TDG/mug DNA glycosylase family. TDG corrects G/T mismatches to G/C pairs. It is capable of hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of the DNA and a mismatched thymine. In addition to the G/T, it can remove thymine also from C/T and T/T mismatches in the order G/T >> C/T > T/T. It has no detectable activity on apyrimidinic sites and does not catalyze the removal of thymine from A/T pairs or from single-stranded DNA. It can also remove uracil and 5-bromouracil from mismatches with guanine. RNF4 interacts with and requires the base excision repair enzymes TDG and APE1 for active demethylation (PMID:20696907). TDG is modified by SUMO-1 and SUMO-2/3. The molecular weight of non-modified TDG is 46 kDa and modified TDG is 55-60 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Jiyu Miao	35414793	Int J Biol Sci	WB
Umut Sahin	24637324	J Cell Biol	WB, IP
Etsuko Shibata	24962565	J Biol Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13370-1-AP (TDG antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-TDG (IP:13370-1-AP, 3ug; Detection:13370-1-AP 1:400) with U-937 cells lysate 3000ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 13370-1-AP (TDG antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



human brain tissue were subjected to SDS PAGE followed by western blot with 13370-1-AP (TDG antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.