For Research Use Only

CD86 Polyclonal antibody

Catalog Number: 13395-1-AP 233 Publications



Basic Information

Applications

Catalog Number: 13395-1-AP Concentration:

650 ug/ml

Source: Rabbit Isotype:

AG4063

Immunogen Catalog Number:

Observed MW: 60-80 kDa

Tested Applications:

WB, IF/ICC, ELISA Cited Applications: WB, IF, Cell treatment

Species Specificity: human, mouse **Cited Species:**

human, mouse, rat, rabbit

GenBank Accession Number:

BC040261 GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000114013 UNIPROT ID: P42081 Full Name:

CD86 molecule Calculated MW: 329 aa, 38 kDa

Positive Controls:

WB: Daudi cells, Raji cells, Ramos cells, RAW 264.7

Purification Method:

WB 1:1000-1:4000 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

IF/ICC: Raji cells,

Background Information

CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of CD86 with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of CD86 with cytotoxic T $lymphocyte-associated\ protein\ 4\ negatively\ regulates\ T-cell\ activation\ and\ diminishes\ the\ immune\ response.\ This$ antibody detects a band of approximately 65-70 kDa, consistent with glycosylated form of CD86 (PMID: 8592066, 21849678).

Notable Publications

Author	Pubmed ID	Journal	Application
Liping Xu	36179453	Tissue Cell	
Qiong Tang	36108436	Ecotoxicol Environ Saf	WB
Bo Zhang	32917854	Cell Death Dis	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13395-1-AP (CD86 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed Raji cells using CD86 antibody (13395-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).