For Research Use Only

CCL19/MIP-3 beta Polyclonal antibody, PBS Only

Antibodies | ELISA kits | Proteins WWW.ptglab.com


Catalog Number:13397-1-PBS


Basic Information	Catalog Number: 13397-1-PBS	GenBank Accession Number: BC027968	Purification Method: Antigen affinity purification
	Concentration: 1 mg/ml	GeneID (NCBI): 6363	
	Source: Rabbit	UNIPROT ID: Q99731	
	Isotype: IgG	Full Name: chemokine (C-C motif) ligand 19	
	Immunogen Catalog Number: AG4200	Calculated MW: 98 aa, 12 kDa	
Applications	Tested Applications: WB, IF/ICC, FC (Intra), Indirect ELISA		
	Species Specificity: human		
Background Information	CCL19, also known as MIP-3 β , is one of several CC cytokine genes clustered on the p-arm of chromosome 9. CCL19 may play a role in normal lymphocyte recirculation and homing. It also plays an important role in trafficking of T cells in thymus, and in T cell and B cell migration to secondary lymphoid organs. It specifically binds to chemokine receptor CCR7.		
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS Only		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





THP-1 cells were subjected to SDS PAGE followed by western blot with 13397-1-AP (MIP-3 Beta antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13397-1-PBS in a different storage buffer formulation. Immunofluorescent analysis of (4% PFA) fixed A549 cells using MIP-3 Beta antibody (13397-1-AP) at dilution of 1:400 and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit1gG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 13397-1-PBS in a different storage buffer formulation.



1X10^6 A549 cells were intracellularly stained with 0.4 ug Anti-Human MIP-3 Beta (13397-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 13397-1-PBS in a different storage buffer formulation.