## For Research Use Only

## OTX2 Polyclonal antibody

Catalog Number: 13497-1-AP

Featured Product

**30 Publications** 



**Basic Information** 

Catalog Number: GenBank Accession Number: 13497-1-AP BC032579 GeneID (NCBI): Concentration: 700 ug/ml 5015

**UNIPROT ID:** Source: Rabbit P32243 Full Name: Isotype:

orthodenticle homeobox 2

Calculated MW: Immunogen Catalog Number: AG4323 297 aa, 32 kDa Observed MW:

31-35 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:16000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200

**Applications** 

**Tested Applications:** 

WB, IHC, FC (Intra), IP, ELISA

**Cited Applications:** WB, IHC, IP, CoIP, ChIP Species Specificity: human, mouse, rat **Cited Species:** 

human, mouse, chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Y79 cells, mouse retina tissue, rat retina tissue

IP: Y79 cells.

IHC: human gliomas tissue,

**Background Information** 

The orthodenticle homeobox 2, encoded by OTX2 gene, is a key transcription factor in developmental processes. In particular, it is required for the early specification of the brain and the embryonic development of sensory organs, including the, pineal gland, pituitary gland, inner part of the ear, eyes, and optic nerve. In later stages, it is important for maintaining intact retina and brain function. In addition, it acts as a transcriptional repressor and a gatekeeper of myogenic and neuronal differentiation in medulloblastoma cells. OTX2 binds to the MyoD1 core enhancer through its homeobox domain and the remarkable repressor activity exhibited by the homeobox domain renders OTX2 transcriptionally repressive

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Kyle S. Smith	36131015	Nature	WB
Meng Li	36273157	J Exp Clin Cancer Res	WB,ChIP
Yue Li	29031852	Neuropharmacology	IHC

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 13497-1-AP (OTX2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas using 13497-1-AP (OTX2 antibody) at dilution of 1:100 (under 40x lens)



IP result of anti-OTX2 (IP:13497-1-AP, 3ug; Detection:13497-1-AP 1:1500) with Y79 cells lysate 2400ug.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human OTX2 (13497-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgC(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).