### For Research Use Only

# TDRKH specific Polyclonal antibody

Isotype:

Catalog Number: 13528-1-AP

Featured Product

18 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 13528-1-AP BC032690
Size: GeneID (NCBI): 550 µg/ml 11022
Source: UNIPROT ID: Rabbit Q9Y2W6

gG tudor and KH domain containing

Immunogen Catalog Number:Calculated MW:AG4422606 aa, 67 kDaObserved MW:

67 kDa

Full Name:

Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

WB 1:2000-1:10000

# **Applications**

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, RIP Species Specificity: human, mouse, rat Cited Species:

human, mouse, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

IF/ICC : HeLa cells,

WB: NCCIT cell, mouse brain tissue

IP: mouse brain tissue,
IHC: human gliomas tissue,

# **Background Information**

Tudor and KH domain-containing protein(TDRKH)Tudor domains are protein modules that mediate protein-protein interactions, potentially by binding to methylated ligands. Both KH and tudor domains are involved in binding RNA or single-strand DNA. Tudor also is a germ cell-specific protein with multiple Tudor domains and is involved in germ plasm formation and germ cell specification. TDRKH is one Tudor protein with a single Tudor domain.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Deqiang Ding	29018194	Nat Commun	IF, WB
Odetta Antico	34767452	Sci Adv	WB
Minjie Tan	32486081	Cells	WB,IHC,IF,IP

# Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13528-1-AP (TDRKH specific antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



NCCIT cell were subjected to SDS PAGE followed by western blot with 13528-1-AP (TDRKH specific antibody at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-TDRKH specific (IP:13528-1-AP, 4ug; Detection:13528-1-AP 1:7000) with mouse brain tissue lysate 4000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 13528-1-AP (TDRKH specific antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).