#### For Research Use Only

# **HMGCR** Polyclonal antibody

Catalog Number: 13533-1-AP

Featured Product 53 Publications

BC033692

3156

GeneID (NCBI):

GenBank Accession Number:



**Basic Information** 

Catalog Number: 13533-1-AP Concentration:

**UNIPROT ID:** Source: Rabbit P04035 Full Name: Isotype: 3-hydroxy-3-methylglutaryl-

Coenzyme A reductase Immunogen Catalog Number: AG4444

Calculated MW:

888 aa, 95 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions: IHC 1:500-1:2000

### **Applications**

**Tested Applications:** 

IHC, ELISA

300 μg/ml

**Cited Applications:** 

IHC, IF

Species Specificity: human, mouse, rat **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

IHC: human liver cancer tissue, human small intestine tissue, human heart tissue

# **Background Information**

HMGCR(3-hydroxy-3-methylglutaryl-coenzyme A reductase) is also named as HMG-CoA reductase and belongs to the HMG-CoA reductase family. This protein exists as a 97-kDa glycoprotein in the endoplasmic reticulum (ER) and is the rate-determining enzyme of the biosynthesis pathway(PMID:15830340). The ectopic expression of either fulllength HMGCR or its novel splice variant promotes dysregulation of the MVA pathway. And the ectopic expression of cHMGCR-FL(60 kDa) and cHMGCR-D13(55 kDa) can be detected in HepG2 cells(PMID:20696928). This protein is also can exsit as a oligomer(PMID:20696928).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Hai-Yan Wang	34588618	Acta Pharmacol Sin	IHC
Wei Xu	32935679	Food Funct	
Liang Wei Wang	31518366	PLoS Pathog	

## Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

# Selected Validation Data



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 13533-1-AP (HMGCR antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 13533-1-AP (HMGCR antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).