

For Research Use Only

YAP1 Polyclonal antibody, PBS Only

Catalog Number: 13584-1-PBS

Featured Product



Basic Information

Catalog Number:

13584-1-PBS

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4510

GenBank Accession Number:

BC038235

GeneID (NCBI):

10413

UNIPROT ID:

P46937

Full Name:

Yes-associated protein 1, 65kDa

Calculated MW:

504 aa, 54 kDa

Observed MW:

65-75 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, IP, ELISA

Species Specificity:

human, mouse, rat

Background Information

Yes-associated protein 1 (YAP1) is a transcriptional regulator which can act both as a coactivator and a corepressor and is the critical downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Plays a key role to control cell proliferation in response to cell contact. Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. The presence of TEAD transcription factors are required for it to stimulate gene expression, cell growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction. Isoform 2 and isoform 3 can activate the C-terminal fragment (CTF) of ERBB4 (isoform 3). Increased expression seen in some liver and prostate cancers (PMID: 31613226, 32488048, 33520338). Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level). It is activated by phosphorylation and degraded by ubiquitination (PMID: 20048001). This antibody is a rabbit polyclonal antibody. The calculated molecular weight of YAP1 is 54 kDa, but routinely observed at 65-75 kDa by Western Blot (PMID: 28230103, 33264286, 36255405).

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13584-1-AP (YAP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 13584-1-AP (YAP1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.




IP result of anti-YAP1 (IP:13584-1-AP, 3ug; Detection:13584-1-AP 1:700) with NIH/3T3 cells lysate 1200ug. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of HepG2 cells using 13584-1-AP (YAP1 antibody) at dilution of 1:50 and Rhodamine-Goat anti-Rabbit IgG. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 13584-1-AP (YAP1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 13584-1-AP (YAP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.






Various lysates were subjected to SDS PAGE followed by western blot with 13584-1-AP (YAP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 13584-1-AP (YAP1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.



WB result of YAP1 antibody (13584-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-YAP1 transfected HepG2 cells. This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human lung cancer tissue using YAP1 antibody (13584-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (4% PFA) fixed human lung cancer tissue using YAP1 antibody (13584-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using YAP1 antibody (13584-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 13584-1-PBS in a different storage buffer formulation.