

For Research Use Only

CD122/IL-2RB Polyclonal antibody

Catalog Number: 13602-1-AP

1 Publications



Basic Information

Catalog Number: 13602-1-AP	GenBank Accession Number: BC025691	Purification Method: Antigen affinity purification
Size: 600 µg/ml	GeneID (NCBI): 3560	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:50-1:500
Source: Rabbit	ENSEMBL Gene ID: ENSG00000100385	
Isotype: IgG	UNIPROT ID: P14784	
Immunogen Catalog Number: AG3929	Full Name: interleukin 2 receptor, beta	
	Calculated MW: 551 aa, 61 kDa	
	Observed MW: 70-85 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB: NK-92 cells, IHC: mouse spleen tissue,
Cited Applications: IF	
Species Specificity: human, mouse	
Cited Species: mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

The interleukin 2 receptor, which is involved in T cell-mediated immune responses, is present in 3 forms with respect to ability to bind interleukin 2. The low affinity form is a monomer of the alpha subunit and is not involved in signal transduction. The intermediate affinity form consists of an alpha/beta subunit heterodimer, while the high affinity form consists of an alpha/beta/gamma subunit heterotrimer. The beta subunit (IL2RB, also known as CD122 and p75) is involved in receptor mediated endocytosis and transduces the mitogenic signals of nterleukin 2.

Notable Publications

Author	Pubmed ID	Journal	Application
Anmar Abu-Romman	37516317	Ocul Surf	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



NK-92 cells were subjected to SDS PAGE followed by western blot with 13602-1-AP (CD122/IL-2RB antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 13602-1-AP (CD122/IL-2RB antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 13602-1-AP (CD122/IL-2RB antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).