

For Research Use Only

TTL Polyclonal antibody

Catalog Number: 13618-1-AP

Featured Product

34 Publications



Basic Information

Catalog Number:

13618-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4526

GenBank Accession Number:

BC036819

GeneID (NCBI):

150465

UNIPROT ID:

Q8NG68

Full Name:

tubulin tyrosine ligase

Calculated MW:

377 aa, 43 kDa

Observed MW:

43-45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:50-1:500

IF-P 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, canine

Positive Controls:

WB : HeLa cells, mouse brain, rat brain

IHC : mouse brain tissue,

IF-P : mouse brain tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Tubulin-tyrosine ligase (TTL) is the enzyme responsible for the reversible addition of a tyrosine residue at the carboxyl end of alpha-tubulin. TTL forms stable complexes with tubulin and inhibit tubulin polymerization. TTL is frequently suppressed during tumor progression with resulting accumulation of dephosphorylated alpha-tubulin in tumor cells. TTL suppression in human cancers is associated with increased tumor aggressiveness.

Notable Publications

Author	Pubmed ID	Journal	Application
Lisa Schulz	36131110	Sci Rep	WB
Shiwei Liu	30232450	Nature	WB
Tianhong Wang	29026081	Nat Commun	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13618-1-AP (TTL antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 13618-1-AP (TTL antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 13618-1-AP (TTL antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of TTL antibody (13618-1-AP; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-TTL transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 13618-1-AP (TTL antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.