For Research Use Only

COMP Polyclonal antibody

Catalog Number:13641-1-AP

Featured Product

3 Publications



Basic Information	Catalog Number: 13641-1-AP	GenBank Accession Number: BC033676	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	, and generating particular
	700 µg/ml	1311	
	Source:	UNIPROT ID:	
	Rabbit	P49747	
	Isotype: IgG	Full Name: cartilage oligomeric matrix protein	
	Immunogen Catalog Number:	Calculated MW:	
	AG4575	757 aa, 83 kDa	
Applications	Tested Applications: ELISA		
	Cited Applications: WB, IF, IHC		
	Species Specificity: human		
	Cited Species:		
	human, mouse, rat		
Background Information	family. COMP is expressed by hy expression is increased during fra structural matrix components. CC	pertrophic chondrocytes and osteoblas	ts during endochondral ossification, and it e assembly of collagen fibrils and other e ECM proteins, including fibronectin,
	family. COMP is expressed by hy expression is increased during fra structural matrix components. CC collagen I, II IX, XII, and XIV, matr	pertrophic chondrocytes and osteoblas acture healing. COMP is involved in the MP interacts with a number of cartilag	ts during endochondral ossification, and it e assembly of collagen fibrils and other re ECM proteins, including fibronectin,
	family. COMP is expressed by hy expression is increased during fra structural matrix components. CC collagen I, II IX, XII, and XIV, matr	pertrophic chondrocytes and osteoblas octure healing. COMP is involved in the MP interacts with a number of cartilag ilins, as well as proteoglycans such as	e ECM proteins, including fibronectin, aggrecan and others.
Background Information	family. COMP is expressed by hy expression is increased during fra structural matrix components. CC collagen I, II IX, XII, and XIV, matr Author Yunjiao Wang	pertrophic chondrocytes and osteoblas acture healing. COMP is involved in the MP interacts with a number of cartilag ilins, as well as proteoglycans such as Pubmed ID Journal	ts during endochondral ossification, and it e assembly of collagen fibrils and other e ECM proteins, including fibronectin, aggrecan and others. Application
	family. COMP is expressed by hy expression is increased during fra structural matrix components. CC collagen I, II IX, XII, and XIV, matr Author Yunjiao Wang Tobias G Schips	pertrophic chondrocytes and osteoblas octure healing. COMP is involved in the MP interacts with a number of cartilag ilins, as well as proteoglycans such as Pubmed ID Journal 31225686 Cell Prolif	ts during endochondral ossification, and it e assembly of collagen fibrils and other re ECM proteins, including fibronectin, aggrecan and others. Application IF

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer. Selected Validation Data