For Research Use Only

LTA4H Polyclonal antibody

Catalog Number: 13662-1-AP

2 Publications



Basic Information

Catalog Number: 13662-1-AP

 Size:
 GeneID (NCBI):

 200 μ g/ml
 4048

Source: UNIPROT ID: Rabbit P09960
Isotype: Full Name:

IgG leukotriene A4 hydrolase

Immunogen Catalog Number:Calculated MW:AG4568611 aa, 69 kDaObserved MW:

57-70 kDa

BC032528

GenBank Accession Number:

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications:

WB. IF

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells, Jurkat cells

IP: HEK-293 cells,

IHC: human skin tissue, human brain tissue, human heart tissue, human kidney tissue, human lung tissue, human ovary tissue, human placenta tissue, human spleen tissue, human testis tissue

Background Information

LTA4H(Leukotriene A-4 hydrolase), also named as LTA4, is essential for the conversion of LTA4 to LTB4, an inflammatory lipid mediator(PMID:10601658). LTA4H also possesses aminopeptidase activity that it can degrade PGP, which is a biomarker for chronic obstructive pulmonary disease and is implicated in neutrophil persistence in the lung(PMID:20813919). This protein has 4 isoforms produced by alternative splicing with the MW of 69, 59, 57, 67 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Wei Gao	31287215	Proteomics	WB
Gan Chen	37122356	JHEP Rep	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data








Immunohistochemical analysis of paraffinembedded human skin using 13662-1-AP (LTA4H antibody) at dilution of 1:100 (under 10x lens).



HEK-293 cells were subjected to SDS PAGE followed by western blot with 13662-1-AP (LTA4H Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-LTA4H (IP:13662-1-AP, 4ug; Detection:13662-1-AP 1:500) with HEK-293 cells lysate 1120ug.