## For Research Use Only

## CPE Polyclonal antibody

Catalog Number: 13710-1-AP

**Featured Product** 

11 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 13710-1-AP BC033866 GeneID (NCBI): Concentration: 153 ug/ml 1363 **UNIPROT ID:** Source: Rabbit P16870

Full Name: Isotype: carboxypeptidase E

Calculated MW: Immunogen Catalog Number: AG4647 476 aa, 53 kDa

> Observed MW: 50 kDa, 52 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF-P 1:200-1:800 IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, IP, ELISA

Cited Applications:

WB, IHC, IF Species Specificity:

human, mouse, rat **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, IP: mouse brain tissue,

IHC: mouse brain tissue, human pituitary tissue, rat

brain tissue

IF-P: mouse brain tissue, IF/ICC: HepG2 cells,

**Background Information** 

CPE(Carboxypeptidase E) is also named as CPH.It belongs to the peptidase M14 family and is involved in the biosynthesis of peptide hormones and neurotransmitters, including insulin. The deduced 476-amino acid human protein contains an N-terminal signal peptide, followed by a 'pro' sequence, a polyarginine stretch, and 2 potential N-glycosylation sites.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Kristen E Rohli	36325514	Function (Oxf)	WB
Jing Yu	34630524	Front Genet	WB,IHC
Seokwon Jo	33115825	Diabetes	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Carboxypeptidase E antibody (13710-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).




mouse brain tissue were subjected to SDS PAGE followed by western blot with 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Carboxypeptidase E antibody (13710-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP result of anti-Carboxypeptidase E (IP:13710-1-AP, 3ug; Detection:13710-1-AP 1:500) with mouse brain tissue lysate 3500ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human pituitary tissue slide using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).