For Research Use Only

GRK3 Polyclonal antibody

Catalog Number:13727-1-AP




Basic Information	Catalog Number: 13727-1-AP	GenBank Accession Number: BC 036797	Purification Method: Antigen affinity purification	
	Size: 500 ug/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG4685	GeneID (NCBI): 157 UNIPROT ID: P35626	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500	
				Full Name: adrenergic, beta, receptor kinase :
		Calculated MW: 688 aa, 80 kDa		
		Observed MW: 70 kDa, 90 kDa		
		Applications	Tested Applications:	Positive C
WB, IHC, ELISA Species Specificity: human, mouse, rat	WB : K-562 cells, HL-60 cells			
	IHC : mous		e brain tissue, human colon tissue	
Note-IHC: suggested antige. TE buffer pH 9.0; (*) Alterno retrieval may be performed buffer pH 6.0	ntively, antigen			
Background Information	GRK3 (G protein coupled receptor kinase 3), also known as ADRBK2 or BARK2, belongs to the G protein coupled receptor kinases (GRKs) family. GRKs are serine/threonine kinases that regulate a large and diverse class of G-protein coupled receptors (GPCRs). GRKs have been shown to play a critical role in the outcome of a variety of physiological and pathophysiological processes including chemotaxis, signaling, migration, inflammatory gene expression and so on (PMID: 28950947, 29483837). GRK3 was identified as a negative regulator of CXCL12/CXCR4 signaling, and has also been shown to be overexpressed in human prostate cancer and found to promote tumor growth and metastasis by enhancing angiogenesis (PMID: 23935208, 24434559). GRK3 has a molecular mass of 75-80 kDa, and also can be ubiquitinated (PMID: 20596523).			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptg W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data






K-562 cells were subjected to SDS PAGE followed by western blot with 13727-1-AP (GRK3 antibody at dilution of 1:2000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13727-1-AP (GRK3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13727-1-AP (GRK3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human normal colon slide using 13727-1-AP (GRK3 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).