### For Research Use Only

# MYST2 Polyclonal antibody

Catalog Number: 13751-1-AP

**Featured Product** 

13 Publications



**Basic Information** 

Catalog Number:

13751-1-AP

BC032640

Size:

GeneID (NCBI):

400 µg/ml

11143

Source:

Rabbit

O95251

Isotype:

GenBank Accession Number:

BC032640

UNCBI):

UNIPROT ID:

O95251

Full Name:

gG MYST histone acetyltransferase 2

Immunogen Catalog Number:Calculated MW:AG4693611 aa, 71 kDaObserved MW:

Observed 83 kDa Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF 1:50-1:500

**Applications** 

Tested Applications: IF/ICC, IHC, IP, WB, ELISA

Cited Applications: WB, IP, IF, IHC, CoIP Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: HeLa cells, human testis tissue, Jurkat cells, NIH/3T3 cells, MCF-7 cells, PC-3 cells

IP: HeLa cells.

IHC : human ovary tumor tissue, human breast cancer

tissue, mouse kidney tissue

IF: NIH/3T3 cells,

## **Background Information**

MYST2, also named as HBO1, HBOa MOZ, YBF2/SAS3, SAS2 and TIP60 protein 2, belongs to the MYST (SAS/MOZ) family. It specifically represses AR mediated transcription. MYST2 is a candidate oncogene. It enhances the anchorage-independent growth of breast cancer cells.(PMID:19372580) MYST2 is a histone acetyltransferase (HAT) which could exert oncogenic function in breast cancer. It is an important downstream molecule of ERa, and ERK1/2 signaling pathway may involved in the expression of HBO1 increased by E2.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Mylinh T Duong	23955388	Cancer Res	WB
Wenbo Liu	36355419	eLife	WB
Wang Wen-zhong WZ	21040551	J Exp Clin Cancer Res	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 13751-1-AP (MYST2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 13751-1-AP (MYST2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human ovary tumor using 13751-1-AP (MYST2 antibody) at dilution of 1:50 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human ovary tumor using 13751-1-AP (MYST2 antibody) at dilution of 1:50 (under 40x lens)



IP result of anti-MYST2 (IP:13751-1-AP, 4ug; Detection:13751-1-AP 1:500) with HeLa cells lysate 2480ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed NIH/3T3 cells using 13751-1-AP (MYST2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).