For Research Use Only

Entactin Polyclonal antibody Catalog Number:13766-1-AP 4 Publications




Basic Information	Catalog Number: 13766-1-AP	GenBank Access BC045606	sion Number:	Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:200-1:800 IF/ICC 1:50-1:500		
	Concentration:	GenelD (NCBI):				
	450 ug/ml Source:	4811 UNIPROT ID:				
	Rabbit	P14543				
	lsotype: IgG	Full Name: nidogen 1				
	Immunogen Catalog Number: AG4749	Calculated MW: 1247 aa, 136 kDa Observed MW: 136-150 kDa				
Applications	Tested Applications: WB, IHC, IF/ICC, IP, ELISA		Positive Co	Positive Controls:		
	Cited Applications: mouse pl WB, IF, CoIP IP : huma Species Specificity:			WB : human placenta tissue, mouse heart tissue, mouse placenta tissue IP : human placenta tissue,		
			IHC : human	an stomach cancer tissue,		
	Cited Species: IF/ICC : HepG2 cells, human, rat					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	Nidogen 1 (Entactin), first identified from a basement membrane-secreting cell line and the murine EHS tumor, is together with perlecan, laminin, and collagen IV, a ubiquitous component of basement membranes (PMID: 19530222). Nidogen-1 is made by mesenchymal cells and is incorporated into the epithelial BM. It is a sulfated glycoprotein of 150 kDa with three globular domains, G1, G2 and G3, and contains binding sites for other ECM molecules (PMID: 10958695, 10671374).					
Notable Publications	Author	Pubmed ID	Journal		Application	
	Pengzhen Cheng	35310379	Bioact Mater		WB,CoIP	
	Emna Ouni	35341935	Matrix Biol		IF	
	Huan Zhang	39472581	Signal Transduct Target Ther		WB	
otorage	Storage: Store at -20°C. Stable for one yes Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -2	nd 50% glycerol pH 7.3	3.			


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13766-1-AP (Entactin antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-Entactin (IP:13766-1-AP, 4ug; Detection:13766-1-AP 1:200) with human placenta tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 13766-1-AP (Entactin antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Entactin antibody (13766-1-AP) at dilution of 1:200 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).