For Research Use Only

NCAM2 Polyclonal antibody Catalog Number:13850-1-AP 4 Publications




Basic Information	Catalog Number: 13850-1-AP	GenBank Accession Number: BC052946		Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilutions: WB 1:5000-1:50000	
	240 ug/ml	4685			
	Source: Rabbit	O15394 protein Full Name: neural cell adhesion molecule 2		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500	
	lsotype: IgG				
	Immunogen Catalog Number: AG4821				
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IP, ELISA WB : mous Cited Applications: tissue		e brain tissue, human brain tissue, rat brai		
	IHC, IF IP : mouse		IP : mouse b	rain tissue,	
	Species Specificity: IHC : mouse brain tissue, human glioma: human, mouse, rat IHC : mouse brain tissue, human glioma:		brain tissue, human gliomas tissue		
	Cited Species: mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	adhesion molecule 2 (NCAM2, al: expressed in the brain, which ma primary olfactory axons or in neu has two isoforms: NCAM2.1 beari	o known as NCAM2 y play important rol rodegenerative dise ng transmembrane a	1) is a single-pass ty es in selective fascic eases such as Alzheir and cytoplasmic dom	ID: 9729879; 20439718). Neural cell pe I membrane protein that is highly ulation and zone-to-zone projection of tl ner's disease (PMID: 26611261). NCAM2 ains and the shorter NCAM2.2 isoform, f NCAM2 is around 125 kDa and 93 kDa	
	adhesion molecule 2 (NCAM2, al expressed in the brain, which ma primary olfactory axons or in neu has two isoforms: NCAM2.1 beari which is GPI-anchored (PMID: 933	o known as NCAM2 y play important rol rodegenerative dise ng transmembrane a	1) is a single-pass ty es in selective fascic eases such as Alzheir and cytoplasmic dom	pe I membrane protein that is highly ulation and zone-to-zone projection of th ner's disease (PMID: 26611261). NCAM2 ains and the shorter NCAM2.2 isoform, f NCAM2 is around 125 kDa and 93 kDa	
	adhesion molecule 2 (NCAM2, al: expressed in the brain, which ma primary olfactory axons or in neu has two isoforms: NCAM2.1 beari which is GPI-anchored (PMID: 933 (PMID: 9334170; 3171685).	to known as NCAM2 y play important rol rodegenerative dise ng transmembrane a 4170; 9221781). The	 is a single-pass ty es in selective fascic eases such as Alzheir and cytoplasmic dom e molecular weight o 	pe I membrane protein that is highly ulation and zone-to-zone projection of tl ner's disease (PMID: 26611261). NCAM2 ains and the shorter NCAM2.2 isoform,	
	adhesion molecule 2 (NCAM2, al: expressed in the brain, which ma primary olfactory axons or in neu has two isoforms: NCAM2.1 beari which is GPI-anchored (PMID: 933 (PMID: 9334170; 3171685).	so known as NCAM2 y play important rol. rodegenerative dise ng transmembrane a 4170; 9221781). The Pubmed ID	1) is a single-pass ty es in selective fascic eases such as Alzheir and cytoplasmic dom e molecular weight o Journal	pe I membrane protein that is highly ulation and zone-to-zone projection of the ner's disease (PMID: 26611261). NCAM2 ains and the shorter NCAM2.2 isoform, f NCAM2 is around 125 kDa and 93 kDa Application	
Background Information	adhesion molecule 2 (NCAM2, al: expressed in the brain, which ma primary olfactory axons or in neu has two isoforms: NCAM2.1 beari which is GPI-anchored (PMID: 933 (PMID: 9334170; 3171685). Author Jia-Qi Ma	so known as NCAM2 y play important rol rodegenerative dise ng transmembrane a 4170; 9221781). The Pubmed ID 39635132	1) is a single-pass ty es in selective fascic eases such as Alzheir and cytoplasmic dom e molecular weight o Journal iScience	pe I membrane protein that is highly ulation and zone-to-zone projection of th ner's disease (PMID: 26611261). NCAM2 ains and the shorter NCAM2.2 isoform, f NCAM2 is around 125 kDa and 93 kDa Mapplication IF	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data






Various lysates were subjected to SDS PAGE followed by western blot with 13850-1-AP (NCAM2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13850-1-AP (NCAM2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13850-1-AP (NCAM2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-NCAM2 (IP:13850-1-AP, 4ug; Detection:13850-1-AP 1:5000) with mouse brain tissue lysate 2100 ug.