

For Research Use Only

NDUFAF2 Polyclonal antibody

Catalog Number: 13891-1-AP **3 Publications**



Basic Information

Catalog Number:

13891-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4853

GenBank Accession Number:

BC033965

GeneID (NCBI):

91942

UNIPROT ID:

Q8N183

Full Name:

NADH dehydrogenase (ubiquinone) 1
alpha subcomplex, assembly factor 2

Calculated MW:

169 aa, 20 kDa

Observed MW:

25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

IHC 1:50-1:500

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

WB, IP

Species Specificity:

human, mouse

Cited Species:

human

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : HepG2 cells,

IP : SH-SY5Y cells,

IHC : mouse skeletal muscle tissue,

Background Information

Mimitin, also named as MMTN, B17.2L, NDUFAF2 and NDUFA12L, is a small mitochondrial protein upregulated by proinflammatory cytokines at the transcriptional and protein levels. It is reported as a possible chaperone for assembly of mitochondrial complex I. The MW of Mimitin is 17-25 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Ostergaard Elsebet E	21617257	J Med Genet	WB
Margherita Protasoni	31912925	EMBO J	WB
Sujin Park	36738241	Redox Rep	IP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 13891-1-AP (NDUFAF2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 13891-1-AP (NDUFAF2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-NDUFAF2 (IP:13891-1-AP, 4ug; Detection:13891-1-AP 1:500) with SH-SY5Y cells lysate 1600ug.