

# NDUFAF2 Polyclonal antibody

Catalog Number: 13891-1-AP

3 Publications

## Basic Information

## Catalog Number:

13891-1-AP

## Size:

800 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG4853

## GenBank Accession Number:

BC033965

## GeneID (NCBI):

91942

## UNIPROT ID:

Q8N183

## Full Name:

NADH dehydrogenase (ubiquinone) 1  
alpha subcomplex, assembly factor 2

## Calculated MW:

169 aa, 20 kDa

## Observed MW:

25 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 µg for 1.0-3.0 mg of total  
protein lysate

IHC 1:50-1:500

## Applications

## Tested Applications:

IHC, IP, WB, ELISA

## Cited Applications:

WB, IP

## Species Specificity:

human, mouse

## Cited Species:

human

**Note-IHC: suggested antigen retrieval with  
TE buffer pH 9.0; (\*) Alternatively, antigen  
retrieval may be performed with citrate  
buffer pH 6.0**

## Positive Controls:

WB : HepG2 cells,

IP : SH-SY5Y cells,

IHC : mouse skeletal muscle tissue,

## Background Information

Mimitin, also named as MMTN, B17.2L, NDUFAF2 and NDUFA12L, is a small mitochondrial protein upregulated by proinflammatory cytokines at the transcriptional and protein levels. It is reported as a possible chaperone for assembly of mitochondrial complex I. The MW of Mimitin is 17-25 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Ostergaard Elsebet E	21617257	J Med Genet	WB
Margherita Protasoni	31912925	EMBO J	WB
Sujin Park	36738241	Redox Rep	IP

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech  
Group brand and is not available to purchase from any  
other manufacturer.**


Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 13891-1-AP (NDUFAF2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 13891-1-AP (NDUFAF2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-NDUFAF2 (IP:13891-1-AP, 4ug; Detection:13891-1-AP 1:500) with SH-SY5Y cells lysate 1600ug.