For Research Use Only

SPAG8 Polyclonal antibody

Catalog Number: 13915-1-AP 2 Publications



Purification Method:

WB 1:500-1:1000

protein lysate

IHC 1:20-1:200

IF/ICC 1:200-1:800

Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Basic Information

Catalog Number:

GenBank Accession Number:

13915-1-AP

BC019247

Concentration:

450 ug/ml

26206

Source:

UNIPROT ID:

Rabbit

Q99932

Isotype:

GenBank Accession Number:

BC019247

UNCBI):

26206

UNIPROT ID:

Rabbit

Q99932

Full Name:

gG sperm associated antigen 8

Immunogen Catalog Number:Calculated MW:AG491045 kDa, 54 kDaObserved MW:

56-70 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC

Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: DU 145 cells,

IP: mouse testis tissue,
IHC: human prostate cancer tissue,

IF/ICC: PC-3 cells,

Background Information

SPAG8, also named as Sperm-associated antigen 8 or HSD-1, is a 426 amino acid protein, which is expressed in testis (germ cells), but not in liver, kidney, prostate and small intestine. SPAG8 may play a role in fertility and microtubule formation through interaction with RANBP9. SPAG8 as testis-specific protein is produced during male germ cell differentiation and functions in a germ cell type- and stage-specific manner during spermatogenesis.

Notable Publications

Author	Pubmed ID	Journal	Application
Gina Esther Merges	35608054	Development	WB
Sili?a Karīna K	21150711	J Immunother	IHC

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human prostate cancer using 13915-1-AP (SPAG8 antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human prostate cancer using 13915-1-AP (SPAG8 antibody) at dilution of 1:100 (under 40x lens)



IP result of anti-SPAG8 (IP:13915-1-AP, 3ug; Detection:13915-1-AP 1:500) with mouse testis tissue lysate 6000ug.



DU 145 cells lysates were subjected to SDS PAGE followed by western blot with 13915-1-AP (SPAG8 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using SPAG8 antibody (13915-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).