## For Research Use Only

## NUP62 Polyclonal antibody

Catalog Number: 13916-1-AP

Featured Product

18 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 13916-1-AP BC014842 GeneID (NCBI): Concentration: 400 ug/ml 23636 **UNIPROT ID:** Source: Rabbit P37198 Full Name: Isotype: nucleoporin 62kDa

Calculated MW: Immunogen Catalog Number: AG4912 522 aa, 53 kDa Observed MW:

62 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

**Tested Applications:** WB, IHC, IP, ELISA Cited Applications: WB, IF, IP

Species Specificity: human, mouse, rat **Cited Species:** 

human, mouse, rat, pig, monkey, chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: PC-3 cells, NIH/3T3 cells

IP: NIH/3T3 cells,

IHC: human stomach tissue.

**Background Information** 

NUP62 belongs to the nucleoporin NSP1/NUP62 family. It is essential component of the nuclear pore complex. The N-terminal is probably involved in nucleocytoplasmic transport. The C-terminal is probably involved in proteinprotein interaction via coiled-coil formation and may function in anchorage of p62 to the pore complex. Defects in NUP62 are the cause of infantile striatonigral degeneration (SNDI). NUP62 consists of a single promoter with a CpG island and three transcribed exons. It is ubiquitously expressed, an observation compatible with its essential role in transporting cargo across the nuclear envelope. (PMID:16029492)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Li Liu	31527085	J Biol Chem	WB
Hao-Long Zeng	29174846	Biochim Biophys Acta	WB
Tianyu Wu	36395215	Science	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



PC-3 cells were subjected to SDS PAGE followed by western blot with 13916-1-AP (NUP62 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



IP result of anti-NUP62 (IP:13916-1-AP, 4ug; Detection:13916-1-AP 1:1000) with NIH/3T3 cells lysate 1200ug.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 13916-1-AP (NUP62 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).