## For Research Use Only

## HORMAD1 Polyclonal antibody

Catalog Number:13917-1-AP 27 Publications




## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 13917-1-AP BC047406 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Concentration: 700 ug/ml 84072 WB 1:500-1:2000 UNIPROT ID: Source: Rabbit Q86X24 Full Name: Isotype: lgG HORMA domain containing 1 Calculated MW: Immunogen Catalog Number: AG4914 45 kDa **Observed MW:** 45 kDa Positive Controls: **Applications Tested Applications:** WB, ELISA WB: HUVEC cells, **Cited Applications:** WB, IF, ColP **Species Specificity:** human, mouse, rat **Cited Species:** human, mouse HORMA domain-containing proteins regulate interactions between homologous chromosomes (homologs) during **Background Information** meiosis in a wide range of eukaryotes [PMID:21079677]. They also implicated in other processes related to crossover formation, including DSB formation, inhibition of promiscuous formation of the synaptonemal complex (SC), and the meiotic prophase checkpoint that monitors both DSB processing and SCs [PMID:19851446]. HORMAD1 first accumulates on the chromosomes during the leptotene to zygotene stages of meiotic prophase I. As germ cells progress into the pachytene stage, HORMAD1 disappears from the synapsed chromosomal regions. However, once the chromosomes desynapse during the diplotene stage, HORMAD1 again accumulates on the chromosome axis of the desynapsed homologs [PMID:19686734]. HORMAD1 exists as various isoform and the range of molecular weight of isoforms are sbout 36-45 kDa. Notable Publications Author Pubmed ID Journal Application PLoS Genet WBIF Yasuhiro Fujiwara 32931493 Mol Biol Cell Hui Tian IF 33175657 Kodai Hirano 35618762 Commun Biol IF Storage: **Storage** Store at -20°C. Stable for one year after shipment. Storage Buffer PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



HUVEC cells were subjected to SDS PAGE followed by western blot with 13917-1-AP (HORMAD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.