## For Research Use Only

# DCX Polyclonal antibody

Catalog Number: 13925-1-AP

26 Publications



**Basic Information** 

Catalog Number: 13925-1-AP

Size: GeneID (NCBI):  $600 \mu \text{ g/ml}$  1641

Source: UNIPROT ID:

Rabbit 043602

Isotype: Full Name:
IgG doublecortin

Immunogen Catalog Number: Calculated MW:
AG4945 402 aa, 45 kDa

Observed MW: 40 kDa

BC027925

GenBank Accession Number:

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:12000 IHC 1:50-1:500 IF 1:50-1:500

**Applications** 

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications:

WB. IF

Species Specificity: human, mouse, rat Cited Species:

human, rat, mouse, hamster

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: SH-SY5Y cells, fetal human brain tissue

IHC: human gliomas tissue,

IF : SH-SY5Y cells,

## **Background Information**

DCX (Doublecortin) is a microtubule-associated protein that is essential for normal brain development. DCX is expressed in migrating neurons throughout the central and peripheral nervous system during embryonic and postnatal development. Anti-DCX antibodies are widely used for detection of neurogenesis. It's also a common neuron marker since its expression is specific to neurons.

### Notable Publications

Author	Pubmed ID	Journal	Application
Li Cheng	36212696	Front Cell Neurosci	IF
Jia Wang	36499306	Int J Mol Sci	WB,IF
Alison A Chomiak	35601919	iScience	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**



Fetal human brain lysates were subjected to SDS PAGE followed by western blot with 13925-1-AP (DCX antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed SH-SY5Y cells using 13925-1-AP (DCX antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 13925-1-AP (DCX antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13925-1-AP (DCX antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).