## For Research Use Only

## GAPDHS Polyclonal antibody

Catalog Number: 13937-1-AP 15 Publications



**Purification Method:** 

WB 1:500-1:3000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: 13937-1-AP

Size:  $600~\mu\,\text{g/ml}$ Source: Rabbit Isotype:

Immunogen Catalog Number:

AG4988

dehydrogenase, spermatogenic Calculated MW:

BC036373

26330

014556 Full Name:

GeneID (NCBI):

**UNIPROT ID:** 

GenBank Accession Number:

glyceraldehyde-3-phosphate

45 kDa Observed MW: 64 kDa

Positive Controls:

WB: mouse testis tissue, rat testis tissue

**Applications** 

**Tested Applications:** WB, ELISA

**Cited Applications:** 

Species Specificity: mouse, rat, human **Cited Species:** 

human, mouse, rat, chicken

**Background Information** 

 ${\sf GAPDHS} (Glyceral dehyde-3-phosphate\ dehydrogen ase,\ test is-specific)\ is\ also\ named\ as\ {\sf GAPD2},\ {\sf GAPDH2},\ {\sf GAPDS}$ and belongs to the glyceraldehyde-3-phosphate dehydrogenase family. It may play an important role in regulating the switch between different pathways for energy production during spermiogenesis and in the spermatozoon. GAPDHS is a promising contraceptive target because it is specific to male germ cells, essential for sperm motility and male fertility, and well suited to pharmacological inhibition(PMID:20828617).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Jingya Gu	33029279	Oxid Med Cell Longev	WB
Zijun Peng	34741939	J Hazard Mater	WB
Guodong Liu	29344176	Oncol Lett	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13937-1-AP (GAPDHS antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 13937-1-AP (GAPDHS antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.