

For Research Use Only

# Siglec-7/CD328 Polyclonal antibody

Catalog Number: 13939-1-AP

3 Publications



## Basic Information

|  |   |  |
|--|---|--|
| <b>Catalog Number:</b><br>13939-1-AP       | <b>GenBank Accession Number:</b><br>BC028150                                | <b>Purification Method:</b><br>Antigen affinity purification         |
| <b>Concentration:</b><br>450 ug/ml         | <b>GeneID (NCBI):</b><br>27036  | <b>Recommended Dilutions:</b><br>WB: 1:500-1:2000<br>IHC: 1:50-1:500 |
| <b>Source:</b><br>Rabbit                   | <b>UNIPROT ID:</b><br>Q9Y286  |  |
| <b>Isotype:</b><br>IgG                     | <b>Full Name:</b><br>sialic acid binding Ig-like lectin 7                   |  |
| <b>Immunogen Catalog Number:</b><br>AG4220 | <b>Calculated MW:</b><br>467 aa, 51 kDa<br><b>Observed MW:</b><br>65-70 kDa |  |

## Applications

|  |   |
|--|---|
| <b>Tested Applications:</b><br>WB, IHC, ELISA  | <b>Positive Controls:</b>   |
| <b>Cited Applications:</b><br>WB, IHC, IF  | <b>WB :</b> human placenta tissue, human liver tissue, U-937 cells, mouse placenta tissue |
| <b>Species Specificity:</b><br>human   | <b>IHC :</b> human tonsillitis tissue, human liver tissue, human placenta tissue          |
| <b>Cited Species:</b><br>human   |   |
| <b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b> |   |

## Background Information

Sialic acid binding Ig-like lectin 7 (Siglec-7), also known as CD328 or p75/AIRM-1, is a member of the Siglec family of glycan-recognition proteins. Siglec-7 is a type-I transmembrane protein consisting of three extracellular immunoglobulin-like domains that comprise an N-terminal V-set domain and two C2-set domains, a transmembrane region and a cytoplasmic tail containing two tyrosine residues embodied in immunoreceptor tyrosine-based inhibition motif-like motifs (PMID: 32322597; 10567377). It is mainly expressed on immune cells, with low levels on granulocytes, intermediate levels on monocytes, and relatively high levels on a major subset of natural killer cells and a minor subset of CD8+ T cells (PMID: 10567377). Siglec-7 is an inhibitory receptor that negatively regulates the function of NK cells and modulates the immune response through the interaction of sialic acid-containing ligands (PMID: 27312286).

## Notable Publications

| Author          | Pubmed ID | Journal       | Application |
|-----------------|-----------|---------------|-------------|
| Kensuke Yamada  | 33240416  | Oncol Lett    | WB,IHC,IF   |
| Jing Zhang      | 33470057  | FEBS Open Bio |             |
| Rebecca Garnham | 38448753  | Commun Biol   | WB,IF       |

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13939-1-AP (Siglec-7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis slide using 13939-1-AP (Siglec-7 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffin-embedded human tonsillitis slide using 13939-1-AP (Siglec-7 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 13939-1-AP (Siglec-7/CD328 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 13939-1-AP (Siglec-7/CD328 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).