For Research Use Only

PDE8A Polyclonal antibody

Catalog Number: 13956-1-AP

5 Publications



Basic Information

Catalog Number: 13956-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

WB 1:500-1:2400

700 μg/ml Source:

5151 **UNIPROT ID:** 060658

BC060762

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Rabbit Isotype:

Full Name:

protein lysate IHC 1:20-1:200

Calculated MW:

phosphodiesterase 8A

Immunogen Catalog Number: AG5069

93 kDa

Observed MW:

93 kDa, 88 kDa

Applications

Tested Applications:

IHC, IP, WB, ELISA

WB: mouse testis tissue, mouse spleen tissue, SH-SY5Y

Cited Applications: WB. IF

Species Specificity:

human, mouse

Cited Species:

IP: mouse testis tissue,

Positive Controls:

human, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

IHC: human colon cancer tissue,

Background Information

PDE8A(high affinity cAMP-specific and IBMX-insensitive 3',5'-cyclic phosphodiesterase 8A) belongs to the cyclic nucleotide phosphodiesterase family and PDE8 subfamily. PDE8A plays a critical role in the modulation of at least one compartment of cAMP and hence PKA activity during $\,\beta$ -adrenergic receptor (β AR) activation in ventricular myocytes(PMID:20353794). The full length protein contains several sites for protein phosphorylation, Nglycosylation, N-myristoylation, and amidation. It has 5 isoforms produced by alternative splicing with the molecular mass of 93 kDa, 88 kDa, 51 kDa and the isoform 4 and 5 have the same molecular mass of 66 kDa. (PMID:11738832).

Notable Publications

Author	Pubmed ID	Journal	Application
Matthew Riopel	32534258	Mol Metab	WB
Annick Bergeron	26724956	Reprod Fertil Dev	
Amel Lounas	31462694	Sci Rep	WB,IF


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse testis tissue were subjected to SDS PAGE followed by western blot with 13956-1-AP (PDE8A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-PDE8A (IP:13956-1-AP, 5ug; Detection:13956-1-AP 1:1000) with mouse testis tissue lysate 8000ug.



Immunohistochemical analysis of paraffinembedded human colon cancer using 13956-1-AP (PDE8A antibody) at dilution of 1:100 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human colon cancer using 13956-1-AP (PDE8A antibody) at dilution of 1:100 (under 40x lens).