#### For Research Use Only

# ACSL1 Polyclonal antibody

Catalog Number: 13989-1-AP

Featured Product

63 Publications



**Basic Information** 

Catalog Number: 13989-1-AP Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG5059

GenBank Accession Number:

BC050073

GeneID (NCBI):
2180

UNIPROT ID:
P33121

Full Name:

acyl-CoA synthetase long-chain

family member 1 Calculated MW: 78 kDa Observed MW: 68 kDa, 78 kDa **Purification Method:** 

Antigen affinity purification

Recommended Dilutions:

WB: 1:1000-1:6000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:200-1:800 IF/ICC: 1:400-1:1600

**Applications** 

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LO2 cells, mouse cerebellum tissue, mouse liver tissue, rat liver tissue

IP: mouse liver tissue.

IHC: human lung squamous cell carcinoma tissue, human heart tissue, human liver cancer tissue

IF/ICC: HepG2 cells,

## **Background Information**

ACSL1(Long-chain-fatty-acid--CoA ligase 1) is also named as FACL1, FACL2, LACS, LACS1, LACS2 and belongs to the ATP-dependent AMP-binding enzyme family. ACSL1 is a 75 kDa protein that is associated peripherally with the plasma membrane(Brian M Wiczer, etc., 2006). ACSL1 is abundantly expressed in tissues, such as liver and brown fat, that metabolize substantial amounts of triglycerides as fuel, and as such, a deficiency in ACSL1 function could have a more profound affect in those cells, resulting in hepatosteatosis and potentially increased very low density lipoprotein production by the liver or decreased thermogenic capacity in brown adipose tissue(PMID:19429676). An anti-rat ACSL1 antibody recognized a band of the predicted 68 kDa in high-speed supernatant from rat liver and in human and murine SMCs, monocyte-derived macrophages, and murine peritoneal macrophages (PMID:17259370). It has 2 isoforms produced by alternative splicing.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yuxiang Sun	31590050	Colloids Surf B Biointerfaces	WB
De Huang	25242319	Cell Rep	WB
Qixue Wang	30279734	Theranostics	IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 13989-1-AP (ACSL1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-ACSL1 (IP:13989-1-AP, 4ug; Detection:13989-1-AP 1:1000) with mouse liver tissue lysate 3040 ug.



Immunohistochemical analysis of paraffinembedded human lung squamous cell carcinoma tissue slide using 13989-1-AP (ACSL1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ACSL1 antibody (13989-1-AP) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).