For Research Use Only

GRK2 Polyclonal antibody

Catalog Number: 13990-1-AP

Featured Product

12 Publications

BC037963

GenBank Accession Number:



Basic Information

Catalog Number: 13990-1-AP

 Size:
 GeneID (NCBI):

 1200 μ g/ml
 156

Source: UNIPROT ID: Rabbit P25098
Isotype: Full Name:

IgG adrenergic, beta, receptor kinase 1

Immunogen Catalog Number:Calculated MW:AG5071689 aa, 80 kDaObserved MW:

80 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

WB 1:500-1:2000

Applications

Tested Applications:

WB, IHC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IP, CoIP Species Specificity:

human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HL-60 cells, Jurkat cells

IP: HL-60 cells,

IHC: human lymphoma tissue,

Background Information

GRK2 (G-protein coupled receptor kinase 2) is also named as ADRBK1 (Beta-adrenergic receptor kinase 1), BARK, and BARK1. G-protein-coupled receptors (GPCRs) transmit extracellular signals across the cell membrane. GPCR kinases (GRKs) desensitize GPCR signals in the cell membrane. GRK2 translocation from the cytoplasm to the cell membrane in response to extracellular signals is critical to the function of the protein, where GRK phosphorylation is an important component of its translocation to the membrane.(PMID:27412951)

Notable Publications

Author	Pubmed ID	Journal	Application
Fan Yang	26347956	Nat Commun	WB
Chen Zhang	25279970	Cancer Biother Radiopharm	WB
Qiyin Zhou	35641493	Nat Commun	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 13990-1-AP (GRK2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 13990-1-AP (GRK2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-GRK2 (IP:13990-1-AP, 4ug; Detection:13990-1-AP 1:1000) with HL-60 cells lysate 3040ug.



1x10^6 HeLa cells were intracellularly stained with 0.4 ug GRK2 Polyclonal antibody (13990-1-AP) and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).