

For Research Use Only

ROD1 Polyclonal antibody

Catalog Number: 14027-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number:

14027-1-AP

Size:

700 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5172

GenBank Accession Number:

BC044585

GeneID (NCBI):

9991

UNIPROT ID:

O95758

Full Name:

ROD1 regulator of differentiation 1 (S. pombe)

Calculated MW:

60 kDa

Observed MW:

55-59 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:12000

IHC 1:20-1:200

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HL-60 cells, mouse thymus tissue, HeLa cells, Jurkat cells, rat thymus tissue

IHC : human lymphoma tissue,

Background Information

ROD1 is a RNA-binding protein that mediates pre-mRNA alternative splicing regulation, it plays a role in the regulation of cell proliferation, differentiation and migration. ROD1 exist some isoforms and also can be modified with the molecular weight range from 55-59 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Qiong Wu	32477006	Cancer Cell Int	WB,IHC
Ya Zhou	30013660	Oncol Lett	WB
Yingji Chen	35016691	Cancer Cell Int	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HL-60 cells were subjected to SDS PAGE followed by western blot with 14027-1-AP (ROD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lymphoma using 14027-1-AP (ROD1 antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human lymphoma using 14027-1-AP (ROD1 antibody) at dilution of 1:100 (under 40x lens).




Various lysates were subjected to SDS PAGE followed by western blot with 14027-1-AP (ROD1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 14027-1-AP (ROD1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 14027-1-AP (ROD1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Various lysates subjected to SDS PAGE followed by western blot with 14027-1-AP (ROD1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



WB result of ROD1 antibody (14027-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ROD1 transfected HeLa cells.