For Research Use Only

CNOT7 Polyclonal antibody Catalog Number: 14102-1-AP Featured Product

Featured Product 8 Publications




Basic Information	Catalog Number: 14102-1-AP	GenBank Accession Numbe BC 060852	r: Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	253 µg/ml	29883	WB 1:500-1:1000	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG5235	UNIPROT ID: Q9UIV1	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
		Full Name: CCR4-NOT transcription complex, subunit 7 Calculated MW: 285 aa, 33 kDa		
				Observed MW: 33-35 kDa
		Applications	Tested Applications:	Positive Controls:
WB, IP, ELISA	WB		WB : mouse testis tissue,	
Cited Applications: WB, IP	IP : mouse testis tissue,			
Species Specificity: human, mouse, rat	ies Specificity:			
Cited Species: human, mouse				
Background Information	Cnot7, Ccr4-Not complex 7, is a component of CCR4-NOT complex and known to be a transcription factor and a modulator of mRNA degradation in yeast. Mammalian Cnot7 interacts with Tob and its family members, which is known as an inhibitor of BMP signaling through interfering with the Smad system and is involved in bone metabolism [PMID:17451368]. It is one of at least 8 subunits that make up the CCR4-Not transcription complex, which is involved in negative regulation of the pol II directed transcription in yeast24 and may be involved ingloba gene regulation by modulation of TATAA-binding protein (TBP) activity [PMID:10880468].			
Notable Publications	Author	Pubmed ID Journal	Application	
	Masashi Yamaji	28297718 Nature	WB	
	Hsin-Hui Hsieh	34090459 J Inflamm	Lond) WB	
	Roy Matkovic	35013187 Nat Comm	un IP	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 14102-1-AP (CNOT7 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.

116kd→ 66kd→

45kd-

 $35kd \rightarrow$

 $25 \text{kd} \rightarrow$

IP result of anti-CNOT7 (IP:14102-1-AP, 4ug; Detection:14102-1-AP 1:300) with mouse testis tissue lysate 4800ug.