

For Research Use Only

IRF9 Polyclonal antibody

Catalog Number: 14167-1-AP

Featured Product

22 Publications



Basic Information

Catalog Number:

14167-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5365

GenBank Accession Number:

BC035716

GeneID (NCBI):

10379

UNIPROT ID:

Q00978

Full Name:

IRF 9

Calculated MW:

393 aa, 44 kDa

Observed MW:

44-48 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:20-1:200

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

ChIP, IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, mouse heart tissue, rat heart tissue, HepG2 cells, MCF-7 cells, HeLa cells, THP-1 cells

IP: mouse heart tissue,

IHC: human cervical cancer tissue,

IF: HepG2 cells,

Background Information

IRF9 also named ISGF3 is a transcription regulatory factor that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with IRF9/ISGF3G to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of IFN stimulated genes, which drive the cell in an antiviral state.

Notable Publications

Author	Pubmed ID	Journal	Application
Yinglu Li	36206767	Mol Cell	WB
Joshua E Burda	35614216	Nature	IHC
Joshua D Jackson	26883073	Mol Cancer Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 14167-1-AP (IRF9 antibody) at dilution of 1:100 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 14167-1-AP (IRF9 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 14167-1-AP (IRF9 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using IRF9 antibody 14167-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP result of anti-IRF9 (IP:14167-1-AP, 4ug; Detection:14167-1-AP 1:500) with mouse heart tissue lysate 3200ug.