

For Research Use Only

# XPR1 Polyclonal antibody

Catalog Number: 14174-1-AP

Featured Product

8 Publications



## Basic Information

Catalog Number:

14174-1-AP

Size:

700 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5373

GenBank Accession Number:

BC041142

GeneID (NCBI):

9213

UNIPROT ID:

Q9UBH6

Full Name:

xenotropic and polytropic retrovirus receptor

Calculated MW:

696 aa, 82 kDa

Observed MW:

82-84 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

## Applications

Tested Applications:

WB, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : Jurkat cells,

## Background Information

XPR1 is a receptor for xenotropic and polytropic murine leukemia retroviruses and a homolog of yeast Syg1 and plant Pi transporter PHO1 (PMID: 20633538). XPR1 has been identified as an atypical G-protein-coupled receptor. Xenotropic or polytropic retrovirus binding to XPR1 can disrupt the cAMP-mediated signaling function of Xpr1, leading to the apoptosis of infected cells (PMID: 22090134). A band of about 82-100 kDa is probably due to abnormal migration of the protein or post-translation modifications. In addition, a band of about 55-72 kDa may be isoforms and fragments.

## Notable Publications

Author	Pubmed ID	Journal	Application
Xingyao Li	32019887	Proc Natl Acad Sci U S A	WB
Huifang Sun	36584480	Biochem Biophys Res Commun	WB
Yoko Akasu-Nagayoshi	35377528	Cancer Sci	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



1x10<sup>6</sup> Jurkat cells were intracellularly stained with 0.25 ug XPR1 Polyclonal antibody (14174-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Jurkat cells were subjected to SDS PAGE followed by western blot with 14174-1-AP (XPR1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 14174-1-AP (XPR1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.