For Research Use Only

NUP153 Polyclonal antibody

Catalog Number: 14189-1-AP 2 Publications



Basic Information

Catalog Number: 14189-1-AP

Concentration: $600 \mu g/ml$

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG5519

154 kDa

GenBank Accession Number:

BC052965 GeneID (NCBI):

9972 **UNIPROT ID:**

P49790 Full Name:

nucleoporin 153kDa Calculated MW:

154 kDa Observed MW: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2400

Purification Method:

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications: WB

Species Specificity: human, mouse, rat **Cited Species:** human, pig

Positive Controls:

WB: K-562 cells, HeLa cells, A431 cells, HepG2 cells,

MCF-7 cells IP: K-562 cells,

Background Information

 $NUP153 is an FG-repeat containing \, NUP, \, contains \, a \, high-affinity \, site \, for \, importin-\beta \, , \, localizes \, on \, the \, nucleoplasmic$ face of the nuclear pore and is required for assembly of the nuclear basket and import of a subset of nuclear proteins [PMID:18845677]. NUP153 also facilitates nuclear export of mRNA and ribonucleoprotein particles and may have additional functions within the nucleus [PMID:16195343]. The 53BP1-NUP153/importin- β pathway as an important aspect of the DDR network add to an emerging evidence of subcellular trafficking as an integral part of genome surveillance [PMID:22075984].

Notable Publications

Author	Pubmed ID	Journal	Application
Mingyang Li	39826728	Int J Biol Macromol	WB
Yunfeng Long	37897200	Biotechnol J	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 14189-1-AP (NUP153 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-NUP153 (IP:14189-1-AP, 4ug; Detection:14189-1-AP 1:500) with K-562 cells lysate 3600ug.



Various lysates were subjected to SDS PAGE followed by western blot with 14189-1-AP (NUP153 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.