

For Research Use Only

Myocilin Polyclonal antibody

Catalog Number: 14238-1-AP **3 Publications**



Basic Information

Catalog Number:

14238-1-AP

Size:

450 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5492

GenBank Accession Number:

BC029261

GeneID (NCBI):

4653

UNIPROT ID:

Q99972

Full Name:

myocilin, trabecular meshwork
inducible glucocorticoid response

Calculated MW:

57 kDa

Observed MW:

66 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

Applications

Tested Applications:

IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB: human heart tissue, human stomach tissue,
mouse skeletal muscle tissue

IP: mouse skeletal muscle tissue,

Background Information

Myocilin (MYOC), also known as TIGR or GLC1A, is a secreted glycoprotein of the olfactomedin family, originally identified in trabecular meshwork cells after prolonged treatment with glucocorticoids, and, independently, in the retina (PMID: 9176893; 9169133). It may participate in the obstruction of fluid outflow in the trabecular meshwork. Defects in MYOC are the cause of primary open angle glaucoma (POAG) (PMID: 21709622). MYOC has a molecular weight of 55-57 kDa, besides, a 66-kDa form, which could be a result of post-translational modifications, has also been reported in some researches (PMID: 9497363; 12150989).

Notable Publications

Author	Pubmed ID	Journal	Application
Miao Wei	36017046	Int J Ophthalmol	WB
Ziyu Zhou	33987224	Ann Transl Med	WB
Xiaochen Wang	37307020	J Ocul Pharmacol Ther	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human heart tissue were subjected to SDS PAGE followed by western blot with 14238-1-AP (Myocilin antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



IP result of anti-Myocilin (IP:14238-1-AP, 4ug; Detection:14238-1-AP 1:500) with mouse skeletal muscle tissue lysate 4000ug.