For Research Use Only

NCAM1/CD56 Polyclonal antibody, PBS Only

proteintech®

Antibodies | ELISA kits | Proteins

www.ptglab.com

Purification Method:

Antigen affinity purification

Catalog Number: 14255-1-PBS

Featured Product

Basic Information

Catalog Number:

14255-1-PBS

Source: Rabbit

Isotype:

Immunogen Catalog Number:

AG5528

GenBank Accession Number:

BC047244

GeneID (NCBI):

4684

ENSEMBL Gene ID: ENSG00000149294

UNIPROT ID:

P13591 Full Name:

neural cell adhesion molecule 1

Calculated MW: 95 kDa

Observed MW:

120 kDa, 140 kDa, 180 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

Species Specificity:

human, mouse, rat, pig

Background Information

Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronection-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isofroms have also been described (PMID: 1856291).

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS only, pH7.3


Selected Validation Data




WB result of NCAM1/CD56 antibody (14255-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NCAM1/CD56 transfected Neuro-2a cells. This data was developed using the same antibody clone with 14255-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 14255-1-AP (NCAM1/CD56 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 14255-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14255-1-AP (NCAM1/CD56 antibody) at dilution of 1:16000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 14255-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 14255-1-AP (NCAM1/CD56 antibody) at dilution of 1:16000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 14255-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse colon tissue using NCAM1/CD56 antibody (14255-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit 1gG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 14255-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using NCAM1/CD56 antibody (14255-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 14255-1-PBS in a different storage buffer formulation.