For Research Use Only

ITPKA Polyclonal antibody

Catalog Number: 14270-1-AP

Featured Product

10 Publications



Purification Method:

WB 1:500-1:3200

protein lysate

IHC 1:100-1:400

IF/ICC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 14270-1-AP BC026331
Size: GeneID (NCBI): 3706
Source: UNIPROT ID: Rabbit P23677

 Isotype:
 Full Name:

 IgG
 inositol 1,4,5-trisphosphate 3-kinase

Immunogen Catalog Number:

AG5588 Calculated MW:

51 kDa Observed MW: 51 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue,
IP: mouse brain tissue,
IHC: human gliomas tissue,
IF/ICC: HepG2 cells,

Background Information

ITPKA(Inositol-trisphosphate 3-kinase A) belongs to the inositol phosphokinase (IPK) family. It accumulates in dendritic spines and seems to be critically involved in synaptic plasticity. The protein possesses two functional activities. In neurons, ITPKA is targeted to F-actin via an N-terminal actin binding domain (amino acids 1-66) and is relevant for long term potentiation and spatial learning(PMID:20022963).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------------|-----------|-------------------------|-------------|
| Windhorst Sabine S | 21792881 | Int J Cancer | WB,IHC |
| Maohua Liu | 29375702 | Exp Ther Med | WB |
| Jingyun Liu | 36558932 | Pharmaceuticals (Basel) | WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 14270-1-AP (ITPKA antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using ITPKA antibody 14270-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP result of anti-ITPKA (IP:14270-1-AP, 4ug; Detection:14270-1-AP 1:400) with mouse brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 14270-1-AP (ITPKA Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 14270-1-AP (ITPKA Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).