For Research Use Only

GLUD1 Polyclonal antibody

Catalog Number: 14299-1-AP

Featured Product

52 Publications



Basic Information

Catalog Number: GenBank Accession Number: 14299-1-AP BC040132
Concentration: GeneID (NCBI): 2746
Source: UNIPROT ID:

Rabbit P00367 Isotype: Full Name:

gG glutamate dehydrogenase 1

Immunogen Catalog Number: Calculated MW:

AG5694 61 kDa

Observed MW: 45-55 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF-P 1:50-1:500

Applications

Tested Applications: WB, IHC, IF-P, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: HeLa cells, mouse lung tissue, HepG2 cells, LO2 cells, PC-3 cells, mouse liver tissue, rat liver tissue

IP: HeLa cells,

IHC: human gliomas tissue, human breast cancer tissue, human liver tissue, mouse liver tissue

IF-P: mouse brain tissue,

Background Information

Human glutamate dehydrogenase (GDH), an enzyme central to the metabolism of glutamate, is known to exist in housekeeping and nerve tissue-specific isoforms encoded by the GLUD1 and GLUD2 genes, respectively. It catalyses the reversible inter-conversion of glutamate to alpha-ketoglutarate and ammonia, thus interconnecting amino acid and carbohydrate metabolism. GLUD1 might contribute to the formation of specific synapses in the hippocampus such as those formed by the projecting neurons of the entorhinal cortex(PMID: 22138648). GLUD1 has a calculated molecular mass of 61 kDa and an apparent molecular mass of 45-55 kDa with the 53aa transit peptide removed.

Notable Publications

Author	Pubmed ID	Journal	Application
Teresa W-M Fan	36150727	J Immunol	
Weiwei Dai	36256480	J Clin Invest	WB
Jessica B Spinelli	29025995	Science	

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 14299-1-AP (GLUD1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 14299-1-AP (GLUD1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 14299-1-AP (GLUD1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 14299-1-AP (GLUD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 14299-1-AP (GLUD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-GLUD1 (IP:14299-1-AP, 4ug; Detection:14299-1-AP 1:1000) with HeLa cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 14299-1-AP (GLUD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using GLUD1 antibody (14299-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).