For Research Use Only

RNF114 Polyclonal antibody

Catalog Number:14338-1-AP




Basic Information	Catalog Number: 14338-1-AP	GenBank Accession Number BC066919	: Purification Method: Antigen affinity purification
	Size: 500 ug/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG5649	GeneID (NCBI): 55905 UNIPROT ID: Q9Y508 Full Name: ring finger protein 114	Recommended Dilutions: WB 1:1000-1:8000
			IHC 1:300-1:1200 IF/ICC 1:50-1:500
		Observed MW: 26 kDa	
		Applications	Tested Applications:
WB, IHC, IF/ICC, ELISA Species Specificity: human			Jurkat cells, MCF-7 cells human ovary cancer tissue,
	IF/ICC		C : BGC-823 cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	RNF114, also known as ZNF228 or ZNF313, is a member of the zinc-binding protein family. RNF114 contains two C2H2 and Ring finger zinc-binding domains and a ubiquitin-interacting motif (UIM). RNF114 functions as an E3 ligase in the ubiquitin-proteasome system and plays an important role in basic cellular processes in different types of cells. RNF114 is expressed widely in many normal tissues, such as the heart, liver, and kidney. RNF114 has 2 isoforms with the molecular mass of 21 and 26 kDa. RNF114 is a novel determinant of psoriasis susceptibility, and its overexpression is a crucial indicator of epithelial inflammation (PMID: 34104941, 35069903, 37523804).		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 14338-1-AP (RNF 114 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed BGC-823 cells using RNF114 antibody (14338-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).

Various lysates were subjected to SDS PAGE followed by western blot with 14338-1-AP (RNF 114 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.