## For Research Use Only

## RBM6 Polyclonal antibody

Catalog Number: 14360-1-AP 3 Publications



**Basic Information** 

Catalog Number:

14360-1-AP

BC046643

Concentration:

700 µg/ml

Source:

Rabbit

P78332

Isotype:

GenBank Accession Number:

BC046643

GeneID (NCBI):

10180

UNIPROT ID:

P78332

Full Name:

gG RNA binding motif protein 6

Immunogen Catalog Number: Calculated MW:
129 kDa
Observed MW:
129 kDa

Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

**Purification Method:** 

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
IHC 1:20-1:200
IF/ICC 1:50-1:500

**Applications** 

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, RIP

Species Specificity:
human, mouse, rat

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: HepG2 cells, mouse spleen tissue

IP: HeLa cells,

IHC: human pancreas cancer tissue,

IF/ICC : HeLa cells,

**Background Information** 

RBM6, also named as DEF3, is a 1123 amino acid protein, which may interact with FAM168B. RBM6 is expressed ubiquitously in adults tissues. RBM6 specifically binds poly(G) RNA homopolymers in vitro. RBM6 exists some isoforms with MV 129 kDa, 69 kDa, and 59 kDa. onversely, fetal thymus and adult kidney express very little RBM6. In the hemopoietic system of mice, the expression of RBM6 is downregulated upon differentiation of progenitor cells into granulocytes but persists during macrophage development.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Baojun Duan	31207145	Proteomics Clin Appl	IHC
Ruoyang Liu	39900560	J Cell Mol Med	WB
Ping Peng	39344314	Front Biosci (Landmark Ed)	RIP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




HepG2 cells were subjected to SDS PAGE followed by western blot with 14360-1-AP (RBM6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer using 14360-1-AP (RBM6 antibody) at dilution of 1:100 (under 40x lens)



IP result of anti-RBM6 (IP:14360-1-AP, 4ug; Detection:14360-1-AP 1:500) with HeLa cells lysate 2000ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 14360-1-AP (RBM6 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).