## For Research Use Only

# GAA Polyclonal antibody

Catalog Number: 14367-1-AP 8 Publications



**Basic Information** 

Catalog Number: 14367-1-AP

Size: 180 µg/ml Source: Rabbit Isotype:

IgG glucosidase, alpha; acid

Immunogen Catalog Number: Calculated MW: AG5690 105 kDa

Observed MW: 70-76 kDa

BC040431

2548

P10253

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

Tested Applications: WB, IP, IHC, ELISA Cited Applications: WB, IHC, IF

Species Specificity: human, mouse Cited Species:

human, mouse, rat, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: DU 145 cells, mouse liver tissue

IP: mouse liver tissue,

IHC: human pancreas cancer tissue, mouse liver

tissue, mouse skeletal muscle tissue

# **Background Information**

GAA(Alpha-1,4-glucosidase) is a lysosomal enzyme involved in the degradation of glycogen within cellular vacuoles. After translation, GAA undergoes proteolytic processing to form two lengths of lysosomal a-glucosidase, and both N-terminal and C-terminal processing occur. The full sequence can be further processed into a mature form. The 105 kDa GAA can be cleaved into 70 kDa or 76 kDa form.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Atsushi Kato	35838176	Org Biomol Chem	IF
Atsushi Kato	35072486	J Med Chem	IF
Jing Wu	28045567	DNA Cell Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 14367-1-AP (GAA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-GAA (IP:14367-1-AP, 4ug; Detection:14367-1-AP 1:600) with mouse liver tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 14367-1-AP (GAA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).